


**Best
Available
Copy**

AD-A285 074



**U.S. ARMY MEDICAL RESEARCH
INSTITUTE OF CHEMICAL DEFENSE**

USAMRICD-TR-94-02

**Development of an In Vitro Model
Assay System for the Evaluation of
the Effects of Toxic Chemicals on
Human Airways**

**James L. Ellis
Margaret G. Filbert**

March 1994

94-31180



Approved for public release; distribution unlimited

94

**U.S. Army Medical Research
Institute of Chemical Defense
Aberdeen Proving Ground, MD 21010-5425**

DTIC QUALITY INSPECTED 3

DISCLAIMER

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Army or the Department of Defense.

DISPOSITION INSTRUCTIONS

Destroy report when no longer needed. Do not return to the originator.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 1994		3. REPORT TYPE AND DATES COVERED Final 1988 to 1993	
4. TITLE AND SUBTITLE Development of an In Vitro Model Assay System for the Evaluation of the Effects of Toxic Chemicals on Human Airways				5. FUNDING NUMBERS 62787A 3M162787A875	
6. AUTHOR(S) Ellis, James L. Filbert, Margaret G.					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: SGRD-UV-YN Aberdeen Proving Ground, MD 21010-5425				8. PERFORMING ORGANIZATION REPORT NUMBER USAMRICD-TR-94-02	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: SGRD-UV-RC Aberdeen Proving Ground, MD 21010-5425				10. SPONSORING/MONITORING AGENCY REPORT NUMBER USAMRICD-TR-94-02	
11. SUPPLEMENTARY NOTES This task was performed under the scientific services agreement issued by Battelle, Columbus Division, under the Scientific Services Program and the auspices of the US Army Research Office, Research Triangle Park, NC 27709-2211					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The ability of the anticholinesterase agent soman to contract human bronchi was examined. Soman (1-2 uM) had variable effects on human bronchi that had not been stimulated with an electric field stimulator (EFS). In bronchi continuously stimulated by EFS (0.5 Hz, 1 ms, 12 V), soman produced contractions in all tissues examined (12 preparations from 9 humans). In tissues stimulated by EFS, the beta-adrenoreceptor agonist isoproterenol produced relaxations that were greater in magnitude than the contractions produced by soman. The duration of the isoproterenol induced relaxations was variable. Of 12 preparations studied, 3 showed no reversal of the relaxation within 120 min, 6 showed a slow reversal with a reversal time of 106 + 6 min and 3 showed rapid reversal with a 50% reversal time of 14 min. In the latter group the duration of the relaxation produced by isoproterenol was doubled (28+ 2 min) by the M2 muscarinic receptor antagonist AFDX 116 (10 uM). These results show that the isolated human bronchus is a useful model for studying the effects of toxic chemical agents such as soman on the airways. The data obtained with isoproterenol suggest that beta-2 agonists may be useful adjuncts for treating the effects of anticholinesterase agents.					
14. SUBJECT TERMS human bronchi, soman, isoproterenol, AFDX-116				15. NUMBER OF PAGES 15	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UNLIMITED		

FOREWORD

This research was carried out under research plan 1-03-15-0000, protocol No. 1-03-87-015-A-443 "Effect of Anticholinesterase Agents on Isolated Human Airway Smooth Muscle," Task Area 875, WBS TechBase, Task NA 1.4, NA 1.6, NA 2.1.1 and NA 2.2.1; JSA requirements S-A-302 and S-A-303. Part of this study was supported by Battelle, Columbus Division, 505 King Avenue, Columbus, OH 43201 under Scientific Services Program Contract No. DAAL03-86-D-0001, Delivery Order No. 1961; TCN 90-088, under the auspices of the US Army Research Office, Research Triangle Park, NC 27709-2211.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

INTRODUCTION

Exposure of personnel to anticholinesterase nerve agents such as soman produces contraction of bronchial smooth muscle, thus increasing airway resistance. Increase in airway resistance leads to wheezing and shortness of breath which in severe cases may be life-threatening (1-5). The contraction of airway smooth muscle by soman and similar agents is due to the accumulation of acetylcholine released from parasympathetic nerves which supply the main neuronal drive in human airways. The accumulation of acetylcholine is due to the inhibition by soman of cholinesterases which would normally inactivate the acetylcholine by hydrolysis.

Because these agents increase the accumulation of acetylcholine at its receptors, the drug of choice in treating nerve agent poisoning has been the muscarinic antagonist atropine (2). Recently, however, investigators have been studying other drugs as possible therapeutic agents. One such class of compounds has been the *beta*-2 adrenergic receptor agonists typified by isoproterenol. These drugs, which have long been used in the treatment of respiratory diseases such as asthma, have also been shown to functionally antagonize cholinergic responses (6). Animal models have shown that isoproterenol will reverse the contraction produced by soman in both canine (7) and guinea pig (8) airways. Differences between the reversal produced by isoproterenol are apparent between canine and guinea pig airways. In canine airways, the isoproterenol relaxation was rapidly reversed and the response showed tachyphylaxis in that subsequent addition of isoproterenol produced no relaxation (7). In contrast, the relaxations produced by isoproterenol were longer lasting in the guinea pig and were reproducible upon further application of isoproterenol (8). Few studies, however, have examined the effects of nerve agents in human isolated airways; thus, it is unknown whether the animal studies represent the human condition. This study, therefore, examined the ability of soman to contract human isolated bronchi *in vitro* and the ability of isoproterenol to relax soman-contracted airways.

METHODS

Human bronchus.

Macroscopically normal, human lung tissue was obtained from organ donors. The average age of the organ donors was 38 ± 4 years; three were male and six were female. The cause of death was primarily head trauma from motor vehicle accidents or gunshot wounds and from cerebrovascular incidents. The organ donor specimens were placed in Viaspan (Dupont Merck Pharmaceuticals, Wilmington, DE) at 4°C and transferred overnight to the laboratory. On arrival at the laboratory, the specimens were placed in a Krebs' solution gassed with 95% oxygen and 5% carbon dioxide at 4°C. Bronchi (5-10 mm i.d.) were trimmed of surrounding parenchyma and blood vessels. The bronchi were cut longitudinally and prepared as transverse strips 4-5 mm wide. The bronchi were suspended between platinum ring electrodes in 10 ml organ baths containing Krebs' solution. The tissues were connected to Grass FT03 force transducers for the measurement of isometric tension which was recorded on a Gould polygraph. The electrodes were connected to a Grass S44 stimulator whose output was passed through a Medlab Stimu-splitter (Fort Collins, CO) for measurement of current across the electrodes and for signal amplification. The current applied in these experiments was 200-300 mA. Preparations were suspended at an initial tension of 2g and washed with fresh buffer every 15 min for a 60 min equilibration period.

After the equilibration period, tissues were either left unstimulated or stimulated continuously (0.5 Hz, 1 ms, 12 V). Soman (1-2 μ M) was then added to the baths. After the soman-induced contraction had reached a plateau, isoproterenol (1 μ M) was added and the tension was observed until it returned to the pre-isoproterenol level or for 120 min. At this point barium chloride (30 mM) was added to maximally contract the tissues. In some tissues which showed rapid reversal of the relaxation to isoproterenol, the ability of AFDX-116 to potentiate the duration of the isoproterenol-induced relaxation was studied. In this set of experiments, two control relaxations to isoproterenol were obtained; AFDX-116 (1 μ M) was then added 5 min before isoproterenol. Once the relaxation to isoproterenol was completely reversed, 10 μ M AFDX-116 was added 5 min before another addition of isoproterenol. When tension had returned to baseline, barium chloride was added to maximally contract the tissues.

Data analysis.

Contractions induced by soman (a) were expressed as a percentage of the maximum contraction produced by barium chloride (c) (Figure 1). Relaxations by isoproterenol (b) are expressed as a percentage of the contraction by soman (a). A percentage greater than 100 therefore indicates that the relaxation to isoproterenol was greater than the contraction by soman. The duration of the relaxation by isoproterenol is calculated as the 50% recovery time, that is the time taken for the relaxation by isoproterenol to become half of the maximal relaxation.

Drugs.

All drugs were obtained from the Sigma Chemical Co (St. Louis, MO) and prepared fresh daily. Soman was obtained from the US Army Chemical Research Development and Engineering Center, Aberdeen Proving Ground, MD 21010-5425.

RESULTS

Figure 1 shows that addition of soman to the tissue chamber produced contracture after a short latency. This is designated as "a." Application of $1\ \mu\text{M}$ isoproterenol produced relaxation of the muscle tension; the magnitude of the isoproterenol-induced relaxation is designated as "b." The maximum contraction obtained with each tissue was determined by the addition of barium chloride ($30\ \text{mM}$) at the end of the experiment. This is designated as "c." Contractions produced by barium chloride were always greater than those obtained in the presence of soman alone. Figure 1 shows that isoproterenol relaxed the tension below the baseline, i.e., before addition of soman. In unstimulated human bronchi, soman ($1\text{--}2\ \mu\text{M}$) produced either no response ($n=3$) or a small contractile response ($n=3$). In those tissues that contracted, the response to soman application averaged $28\pm 5\%$ of the maximum response produced by barium chloride. The response to barium chloride averaged $1.80\pm 0.4\ \text{g}$ in these tissues. In the three preparations that contracted upon exposure to soman, isoproterenol ($1\ \mu\text{M}$) relaxed the tension below that of the pre-soman tension (Figure 2A). These relaxations were $183\pm 26\%$ of the tension produced by soman. The 50% recovery time for the isoproterenol-induced relaxation was $77\pm 21\ \text{min}$ in these tissues.

In tissues continuously stimulated by EFS ($0.5\ \text{Hz}$, $1\ \text{ms}$, $12\ \text{V}$), 8 out of the 12 preparations showed no contraction, whereas 4 showed a small contraction that subsided over time. The strength of this contraction at its peak averaged $6\pm 2\%$ of the maximum response produced by barium chloride. Despite the observation that the majority of tissues continuously stimulated failed to show a contraction to EFS, they all subsequently contracted in the presence of soman (Figures 2 & 3), and isoproterenol ($1\ \mu\text{M}$) elicited relaxations in all of these tissues. However, the duration of the isoproterenol relaxation was quite variable. Three types of responses to isoproterenol were noted: (1) relaxation with no reversal for 120 min ($n=3$) (Figure 2B); (2) relaxation with slow reversal, having a 50% recovery time of $106\pm 12\ \text{min}$ ($n=6$) (Figure 2C); (3) rapid reversal with 50% recovery time of $14\pm 1\ \text{min}$ ($n=3$) (Figure 3).

In tissues that did not exhibit reversal of relaxation, soman-induced contractures averaged $29\pm 7\%$ of the response to barium chloride. The response to barium chloride in these tissues was $0.8\pm 0.28\ \text{g}$. Relaxations produced by isoproterenol averaged $214\pm 38\%$ of the soman contractions. In tissues showing slow reversal of relaxation, the contracture induced by soman averaged $39\pm 7\%$ of the response to barium chloride. The response to barium chloride in these tissues was $1.87\pm 0.04\ \text{g}$, and isoproterenol produced relaxations averaging $160\pm 14\%$ of the tension produced by soman. In tissues exhibiting rapid reversal of the isoproterenol-induced relaxation, the contraction produced by soman averaged $39\pm 5\%$ of the response to barium chloride. The response to barium chloride in these tissues was $1.83\pm 0.2\ \text{g}$, and isoproterenol produced relaxations averaging $198\pm 7\%$ of the soman-induced contractions.

The effect of AFDX-116, a muscarinic M_2 receptor subtype antagonist, on the duration of the relaxation produced by isoproterenol was examined in the group exhibiting rapid reversal of the isoproterenol-induced relaxation. A initial application of isoproterenol ($1\ \mu\text{M}$) had a 50% recovery time of $14\pm 1\ \text{min}$, and a second application of isoproterenol ($1\ \mu\text{M}$) exhibited a 50% recovery time of $13\pm 1\ \text{min}$. AFDX-116 ($1\ \mu\text{M}$) had little effect on the 50% recovery time for a third application of isoproterenol ($15\pm 2\ \text{min}$), whereas AFDX-116 ($10\ \mu\text{M}$) produced a doubling of the 50% recovery time for a fourth application of isoproterenol ($28\pm 2\ \text{min}$) (Figure 3). In 2 out of the 3 preparations, AFDX-116 ($10\ \mu\text{M}$) produced a small relaxation (8% of that produced by isoproterenol), suggesting that at this concentration AFDX-116 had an effect on both M_3 and M_2 muscarinic receptor subtypes.

DISCUSSION

The results of the present study show that soman is capable of contracting human isolated bronchi. This is in agreement with animal studies where soman has been shown to contract the isolated airways from several species including canine (7), guinea pig (9) and rat (10). The results with unstimulated tissues indicate that the release of acetylcholine in these tissues is variable and is in agreement with studies which have shown that atropine has little effect on intrinsic tone in human airways (11). Soman, however, contracted all the tissues that were continuously stimulated at low frequency (0.5 Hz), even those tissues that showed no contractile response to this stimulation in the absence of soman. It would, therefore, seem prudent that this procedure be used in subsequent studies in human bronchi where the effect of soman and similar agents is to be studied.

The *beta*-2 adrenergic receptor agonist isoproterenol was able to relax all the tissues which exhibited contractions by soman. In each case, the relaxation by isoproterenol was greater than the contraction by soman, indicating that isoproterenol was able to functionally antagonize both the cholinergic response due to soman and the noncholinergic intrinsic tone which is pronounced in human isolated bronchi (11). The relaxations produced by isoproterenol were on the whole long-lasting: in 9 out of 12 preparations studied the 50% recovery time was greater than 60 min. In this respect, the response to isoproterenol in human airways is more like the guinea pig airways which show slower recovery of the isoproterenol relaxant response than do canine airways (7). In 3 out of 12 tissues, the reversal of the isoproterenol response was fairly rapid although still slower than that observed in canine airways. In the latter group, the addition of another dose of isoproterenol produced a relaxation similar to the first relaxation. Although these findings are similar to those observed in the guinea pig, they differ markedly from the results obtained in the canine airway where the subsequent addition of isoproterenol had no effect (7). Also in agreement with the study on guinea pig airways, the M2 muscarinic receptor antagonist AFDX-116 prolonged the relaxation elicited by isoproterenol.

The reversal of the isoproterenol relaxation may be due to a number of factors:

1. Tachyphylaxis. *Beta*-2 adrenoceptors have been shown to be desensitized by repeated administration of an agonist. The continued presence of isoproterenol in the bath may therefore produce internalization of receptors or other events which produce tachyphylaxis. In the present study this is unlikely to be the case, as repeated administration of isoproterenol produced similar results.
2. Breakdown of the isoproterenol. Isoproterenol can be rapidly reduced, leading to loss of activity. This would not explain, however, how the duration of the response to isoproterenol is enhanced by AFDX-116.
3. Functional antagonism. Isoproterenol produces relaxations of airway smooth muscle by activating adenylate cyclase leading to increased cyclic AMP, whereas stimulation of muscarinic receptors produces the opposite effect and will thus tend to oppose relaxations elicited by isoproterenol (12-14). The ability of AFDX-116 to prolong relaxations to isoproterenol suggests that M2 receptors may be involved in this functional antagonism. Supporting this argument is the finding of Fernandes *et al.* (15) which showed that isoproterenol-induced relaxations of methacholine-induced tone in canine trachea were augmented by AFDX-116.

CONCLUSIONS

The human isolated bronchus serves as a very useful model system to study the effect of nerve agents such as soman. Moreover, *beta*-2 adrenergic agonists may serve as useful adjuncts to conventional atropine therapy in the treatment of nerve agent poisoning; furthermore, the usefulness of *beta*-2 agonists may be enhanced by M2 antagonists. Additional studies should be carried out on human airways to examine further possibilities for treatment of nerve agent poisoning. These might include evaluation of new longer acting *beta*-agonists, calcium channel blockers and potassium channel activators.




Figure1. Diagram of research protocol. Soman was added to either unstimulated or stimulated (0.5 Hz, 1 ms, 12 V continuously) human bronchus. The contraction by soman is designated "a". Once the contraction by soman has reached a plateau 1 μ M isoproterenol was added. The relaxation to isoproterenol is designated "b". At the end of the experiment tissues were maximally contracted to 30 mM barium chloride. This maximum contraction is designated "c".




Figure 2. Representative tracings of responses to soman and subsequent relaxations to isoproterenol in human isolated bronchi. A) Response to soman in an unstimulated tissue and subsequent relaxation to soman. B). Response to soman in a continuously stimulated tissue (0.5 Hz, 1 ms, 12 V) in which the relaxation to isoproterenol was maintained for over 120 min. C). Response to soman in a continuously stimulated tissue (0.5 Hz, 1 ms, 12 V) in which the relaxation to isoproterenol returned to baseline.




Figure 3. Representative tracing of contraction by soman in a continuously stimulated (0.5 Hz, 1 ms, 12 V) human isolated bronchus in which the relaxation to isoproterenol was rapidly reversed. AFDX-116 (10 μ M) caused the relaxation to isoproterenol to be doubled in duration compared to the two previous control additions of isoproterenol.

REFERENCES

1. G.B. Koelle and A. Gilman, Anticholinesterase drugs, *Pharmacol Rev.*, 1, 166-216 (1949).
2. F.R. Sidell, Clinical Overview of Organophosphorus Intoxication, in "Proceedings of the Symposium on Respiratory Care of Chemical Casualties." USAMRICD, P-27-84, 1983, pp.15-36.
3. V. Vojvodic, "Toxicology of War Gases", 1st ed., Federal Ministry for National Defense and Health Administration, Military Publishing House. Belgrade, 1981, pp226-228.
4. H. Cullumbine, Actions at Autonomic Effector Sites, in "Cholinesterases and Anticholinesterases," *Handbuch der experimentellen pharmakologic.* (G. Koelle, ed.), Springer-Verlag, Berlin, 1963, p. 510.
5. B. Holmstead, Pharmacology of organophosphorus cholinesterase inhibitors, *Pharmacol. Rev.*, 11, 567-688 (1959).
6. K.G. Rhoden, L.S. Meldrum and P.J. Barnes, Inhibition of cholinergic neurotransmission in human airways by β -2 adrenoceptors, *J. Appl. Physiol.*, 67, 16-22 (1988).
7. M.G. Filbert, D.H. Moore and M. Adler, Relaxation of soman-induced contracture of airway smooth muscle *in vitro*, *Drug Chemical Toxicol.*, 15, 203-215 (1992).
8. M.G. Filbert and G.K. Adams III, Contribution of muscarinic M2 receptors to airway smooth muscle responses., *The Pharmacologist*, 33, 213 (1991).
9. P. Aas, P. Walday, R. Tansø and F. Fonnum, The effect of acetylcholinesterase inhibition on the tonus of guinea pig bronchial smooth muscle, *Biochem. Pharmacol.*, 37, 4211-4216 (1988).
10. P. Walday, E. Fyllingen and P. Aas, Effects of calcium antagonists (w-conotoxin, verapamil, gallopamil, diltiazem) on bronchial smooth muscle contractions induced by soman, *Naunyn-Schmiedeberg's Arch. Pharmacol.*, in press.
11. J.L. Ellis and B.J. Udem, Role of cysteinyl-leukotrienes and histamine in mediating intrinsic tone in human isolated bronchi. *Am. Rev. Respir. Dis.*, in press.
12. G.F. Van den Brink, The model of functional interaction II. Experimental verification of a new model: the antagonism of β -adrenoceptor stimulants and other agonists. *Eur. J. Pharmacol.*, 22, 279-286 (1973).
13. T.J. Torphy, G.A. Rinard, M.G. Rietow and S.E. Mayer, Functional antagonism in canine tracheal smooth muscle: inhibition by methacholine of the mechanical and biochemical responses to isoproterenol, *J. Pharmacol. Exp. Ther.*, 227, 694-699 (1983).
14. T.J. Torphy, S. Zheng, S.M. Peterson, R.R. Fiscus, G.A. Rinard and S.E. Mayer, Inhibitor effect of methacholine on drug-induced relaxation, cyclic AMP accumulation, and cyclic AMP-dependent protein kinase activation in canine tracheal smooth muscle, *J. Pharmacol. Exp. Ther.*, 233, 409-417 (1985).
15. L.B. Fernandes, A.D. Fryer and C.A. Hirshman, M2 muscarinic receptors inhibit isoproterenol-induced relaxation of canine airway smooth muscle, *J. Pharmacol. Exp. Ther.*, 262, 119-126 (1992).

Distribution List

Addresses	Copies	Addresses	Copies
DEFENSE TECHNICAL INFORMATION CENTER ATTN DTIC DDAC CAMERON STATION BUILDING 5 ALEXANDRIA VA 22304-6145	2	DIRECTOR ARMED FORCES MEDICAL INTELLIGENCE CENTER FORT DETRICK MD 21702-5004	1
COMMANDER US ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND FORT DETRICK MD 21702-5012	2	COMMANDER US ARMY INSTITUTE OF DENTAL RESEARCH BUILDING 40 WASHINGTON DC 20307-5300	1
HQDA DASG HCD WASHINGTON DC 20310	1	COMMANDER US ARMY INSTITUTE OF SURGICAL RESEARCH BUILDING 2653 FORT SAM HOUSTON TX 78234-6200	1
DIRECTOR WALTER REED ARMY INSTITUTE OF RESEARCH BUILDING 40 WASHINGTON DC 20307-5100	1	COMMANDANT ACADEMY OF HEALTH SCIENCES US ARMY ATTN HSHA CDC FORT SAM HOUSTON TX 78234-6100	1
COMMANDER US ARMY AEROMEDICAL RESEARCH LABORATORY ATTN SCIENTIFIC INFORMATION CENTER PO BOX 577 FORT RUCKER AL 36362-5000	1	COMMANDANT ACADEMY OF HEALTH SCIENCES US ARMY ATTN HSHA CDM FORT SAM HOUSTON TX 78234-6100	1
COMMANDER US ARMY MEDICAL RESEARCH INSTITUTE OF INFECTIOUS DISEASES BUILDING 1425 FORT DETRICK MD 21702-5011	1	Dr. JOSEPH OSTERMAN DIRECTOR ENVIRONMENTAL AND LIFE SCIENCES OFFICE OF THE DEPUTY DIRECTOR FOR RESEARCH AND ENGINEERING ROOM 3D129 WASHINGTON DC 20301-2300	1
COMMANDER US ARMY RESEARCH INSTITUTE OF ENVIRONMENTAL MEDICINE BUILDING 42 NATICK MA 01760-5007	1	COMMANDER US ARMY TRAINING AND DOCTRINE COMMAND ATTN ATMD FORT MONROE VA 23651	1
COMMANDANT US ARMY CHEMICAL SCHOOL ATTN ATZN CM C FORT MCCLELLAN AL 36205	1		

COMMANDER US ARMY NUCLEAR AND CHEMICAL AGENCY 7500 BACKLICK ROAD BUILDING 2073 SPRINGFIELD VA 22150-3198	1	AFOSR/NL BUILDING RM A217 BOLLING AFB DC 20332	1
BIOLOGICAL SCIENCE DIVISION OFFICE OF NAVAL RESEARCH ARLINGTON VA 22217	1	COMMANDER US ARMY CHEMICAL BIOLOGICAL DEFENSE AGENCY ATTN AMSCB CI ABERDEEN PROVING GROUND MD 21010-5423	1
EXECUTIVE OFFICER NAVAL MEDICAL RESEARCH INSTITUTE NAVAL MEDICINE COMMAND NATIONAL CAPITAL REGION BETHESDA MD 20814	1	LTC DON W KORTE JR BATTELLE MEMORIAL INSTITUTE JM 3 505 KING AVENUE COL UMBUS OH 43201-2695	1
USAF ARMSTRONG LABORATORY/CFTO SUSTAINED OPERATIONS BRANCH BROOKS AFB TX 78235-5000	1	COMMANDER US ARMY MEDICAL RESEARCH INSTITUTE OF CHEMICAL DEFENSE ATTN SGRD UV ZA SGRD UV ZB SGRD UV ZS SGRD UV RC (5 copies) SGRD UV R (11 copies) SGRD UV AI W SGRD UV D SGRD UV P SGRD UV V SGRD UV Y ABERDEEN PROVING GROUND MD 21010-5425	24
DEPARTMENT OF HEALTH AND HUMAN SERVICES NATIONAL INSTITUTES OF HEALTH THE NATIONAL LIBRARY OF MEDICINE SERIAL RECORDS SECTION 8600 ROCKVILLE PIKE BETHESDA MD 20894	1		
STEMSON LIBRARY ACADEMY OF HEALTH SCIENCES BUILDING 2840 RM 106 FORT SAM HOUSTON TX 78234-6100	1		
US ARMY RESEARCH OFFICE ATTN CHEMICAL AND BIOLOGICAL SCIENCES DIVISION PO BOX 12211 RESEARCH TRIANGLE PARK NC 27709-2211	1		